

Impaired social interaction

Nr.	Nursing Diagnosis	Date	Nursing Intervention/s and tasks	Sign.
	<p><b>Impaired social interaction</b></p> <p><b>Related factors:</b></p> <ul style="list-style-type: none"> <li>Absence of available significant others or peers</li> <li>Altered thought processes</li> <li>Communication barriers</li> <li>Environmental barriers</li> <li>Knowledge/skill deficit about ways to enhance mutuality</li> <li>Limited physical mobility</li> <li>Self concept disturbance</li> <li>Sociocultural dissonance</li> <li>Therapeutic isolation</li> </ul> <p><b>Symptoms / Risk factors:</b></p> <ul style="list-style-type: none"> <li>Dysfunctional interaction with others</li> <li>Dysfunctional interaction with peers, family and/or others</li> <li>Family report of change of style or pattern of interaction</li> <li>Observed use of unsuccessful social interaction behaviors</li> <li>Verbalized or observed discomfort in social situations</li> <li>Verbalized or observed inability to receive or communicate a</li> </ul> <p><b>Outcome / Goal:</b></p> <ul style="list-style-type: none"> <li>Patient and family will participate in care and therapies</li> <li>Patient engages in social interaction with significant other</li> </ul>		<p><b>Socialization Enhancement</b></p> <ul style="list-style-type: none"> <li>Ascertain client's perception of situation</li> <li>Be alert to verbal and nonverbal cues including withdrawal, despair and aloneless</li> <li>Encourage enhanced involvement in already established relationships</li> <li>Explore strengths and weaknesses of current network of relationships</li> <li>Give positive feedback when patient reaches out to others</li> <li>Identify support systems available to client</li> </ul>	