

Risk for altered parenting

Nr.	Nursing Diagnosis	Date	Nursing Intervention/s and tasks	Sign.
	<p><b>Risk for altered parenting</b></p> <p><b>Related factors:</b></p> <ul style="list-style-type: none"> <li>Complicated illness</li> <li>Parents</li> <li>Stay in neonatal intensive care unit</li> </ul> <p><b>Symptoms / Risk factors:</b></p> <ul style="list-style-type: none"> <li>Inexperience with caregiving</li> <li>Verbalization of role inadequacy</li> </ul> <p><b>Outcome / Goal:</b></p> <ul style="list-style-type: none"> <li>Demonstrates correct applications and opera. of apnea monitor</li> <li>Parents openly express feelings.</li> <li>Parents will demonstrate necessary skills for caring</li> <li>Verbalized readiness to deal with apneic episodes</li> </ul>		<p><b>Role Enhancement</b></p> <ul style="list-style-type: none"> <li>Allow for verbalization of feelings</li> <li>Allow parents to watch nurse care for infant</li> <li>Encourage parents to participate in care</li> <li>Encourage and promote an opportunity for the mother to hold her newborn</li> <li>Encourage breastfeeding for mother who desires to breastfeed if possible</li> <li>Encourage father to hold infant</li> <li>Explain all procedures and the associated discomforts</li> <li>Give family as much time they need together, with minimum interruption from s</li> <li>Inform parents of neonate's immediate needs</li> <li>Instruct Parents about the use of Apnea monitor</li> <li>Observe the parents interacting with the infant</li> <li>Provide written and/ pictoral instruction regarding the Apnea Monitor</li> <li>Provide praise and support for parents as they learn to use the Apnea Monitor</li> <li>Unite mother, infant ,and family as soon as condition of infant is stable</li> <li>Support each parent's strengths</li> </ul>	