

Risk for peripheral neurovascular dysfu.

Nr.	Nursing Diagnosis	Date	Nursing Intervention/s and tasks	Sign.
	<p>Risk for peripheral neurovascular dysfuncti</p> <p>Related factors: burns of extremities interruption of arterial or venous blood flow Mechanical compression Orthopedic surgery</p> <p>Symptoms / Risk factors: Necrosis of skin Immobilization Edema Paralysis Paresthesia Weak or no peripheral pulses palpable</p> <p>Outcome / Goal: Color of skin normal Patient will not exhibit symptoms of neurovascular compromis peripheral pulses palpable</p>		<p>Circulatory Care: Arterial Insufficiency Administer intravenous (IV) fluids and products as indicated. Apply direct pressure to bleeding site if hemorrhage occurs Assess color, sensation, movement, capillary refill Elevate affected extremities, as appropriate Encourage and assist with early ambulation Ensure that stabilizing devices such as abduction pillow are in correct positio Inspect dressings and drainage device Maintain fluid replacement per protocol Monitor lab. Hgb/Hct and Coagulation studies Monitor vital signs Palpate peripheral pulses noting strength and equality Periodic assessments:sensation, movement, pulse, skin color, and temp.</p> <p>Peripheral Sensation Management Monitor and record if following occur: Change in sensation, Change in movement ability Increasing pain or pain not controlled by medication Pale,mottled, or cyanotic skin Slowed capillary refill more than 3 seconds</p>	